

RHINOPLASTY

DR. JASON CLAIN

POST-OPERATIVE INSTRUCTIONS RHINOPLASTY

CONTACT DR. CLAIN

Office: 914-219-8332

Secretary: (516) 739-2322

Dr. Clain's email:

JClain@nyplasticsurgical.com

Before Surgery

- Do not take any ibuprofen containing products for 2 weeks prior to surgery. Acetaminophen/Tylenol is safe.
- Wear comfy clothes, wear a shirt that buttons down the front so it is easy to get on/off

After Surgery

- Keep your head elevated on pillows, be careful bending down.
- Icing the cheeks can be helpful, do not put pressure on the face
- Gentle movements of the head/neck to prevent bleeding
- Showers are allowed, no swimming/baths/sauna/steam

Activity

- Do not lift anything >5 pounds.
- Keep your head up, do not bend down
- No makeup for 2 weeks. No heavy exercise (straining/sweating) for 6 weeks.

Medications

- Take acetaminophen and ibuprofen for pain. Take the oxycodone for severe pain
- Use the
- You do not need any additional antibiotics.

What To Expect While Healing

- Bruising is normal, this will be better in 1 week and usually gone by 2-3 weeks

Follow-Up

- You will see Dr. Clain 1 week and 2 weeks after surgery. If you do not have an appointment, call his secretary
- It will take 6-12 months to see your final result

RHINOPLASTY

DR. JASON CLAIN